arXiv:astro-ph/0311363v2 20 Apr 2004

Dominant gamma-ray bursts production in the early universe

J. R. Lin[1], S. N. Zhang[1,2,3,4] and T. P. Li[1,4]

[1] *Tsinghua Center for Astrophysics, Tsinghua University, Beijing, 100084, China*

[2] *Physics Department, University of Alabama in Huntsville, Huntsville, AL35899, USA*

[3] *Space Science Laboratory, NASA Marshall Space Flight Center, SD50, Huntsville, AL35812, USA*

[4] *Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, China*

ABSTRACT

It has been known that at least some of the observed gamma-ray bursts (GRBs) are produced at cosmological distances and the GRB production rate may follow the star formation rate. We model the BATSE detected intensity distribution of long GRBs in order to determine their space density distribution and opening angle distribution. Our main results are: the lower and upper distance limits to the GRB production are $z \approx 0.24$ and $z > 10$, respectively; the GRB opening angle follows exponential distribution and the mean opening angle is about 0.03 radian; the peak luminosity appears a better standard candle than the total energy of a GRB.

Subject headings: gamma rays: bursts – stars: formation – cosmology: early universe, reionization

1. Introduction

Recent research results indicate that radiation from gamma-ray bursts (GRBs) may be constrained mostly in narrow beams (Waxman *et al.* 1998; Fruchter *et al.* 1999; Woosley 2001), which suggests that GRBs may be standard candles at cosmological distances (Frail *et al.* 2001). Previous studies of the GRB space density distribution suggested that GRB production rate may follow the observed star formation rate for small values of redshift (Wijers *et al.* 1998), but may increase monotonically for very high values of redshift (Schaefer *et al.* 2001; Norris 2002); however neither the lower limit nor the upper limit of redshift for GRB production could be determined reliably previously because these studies are either limited by statistics or have to rely on some empirical relationships between some statistical properties of selected samples of

GRBs. Because both the space density distribution and beaming effect play important roles for the observed GRB intensity distribution, here we make use of the BATSE/CGRO GRB intensity distribution, which is the largest sample of GRBs collected so far, to determine directly the GRB space density and opening angle distribution.

2. Models for GRB intensity distribution

For the GRB space density distribution, we assume the following model, under the cosmological parameters of $\Omega = 1$, $H_0 = 50$ km s^{-1} Mpc^{-1} and $q_0 = 0.5$, in order to compare with previous star formation rate measurements (Steidel *et al.* 1999).

$$n(1+z) = \begin{cases} 0, & \text{if}(z < z_0) \\ n_0(1+z)^{3.5}, & \text{if}(z_0 \le z < z_{break}) \\ n_0(1+z_{break})^{3.5}, & \text{if}(z \ge z_{break}) \end{cases} . \tag{1}$$

The functional form is taken from the measured star formation rate (Steidel *et al.* 1999) with $z_0 > 0.2$ and $z_{break} \approx 1$. Our main goal in this paper is to determine z_0 and z_{break} for GRBs from the observed GRB intensity distribution, in order to answer the question where in the universe GRBs are produced predominantly.

For the GRB opening angle distribution, we study three cases when the GRB opening angle θ follows Gaussian, exponential, or power-law distributions. For a Gaussian distribution of θ, the probability that the opening angle equals to a given angle is,

$$P(\theta) = \frac{1}{\sqrt{2\pi}\sigma} \mathrm{e}^{-\frac{(\theta-\theta_0)^2}{2\sigma^2}}. \tag{2}$$

We assume that the total energy E_0 is the same for for every long GRB, i.e., they can be used as standard candles(Frail *et al.* 2001; Balazs *et al.* 2003) and the opening angle θ is very small, the probability that the observed fluence f is higher than a specified value F is then,

$$\begin{aligned} P(f \ge F) &= P(\Omega \approx \frac{\pi}{4}\theta^2 \le \frac{E_0(1+z)}{F d_L^2}) \\ &= \int_0^{(\frac{4E_0(1+z)}{\pi F d_L^2})^{\frac{1}{2}}} \frac{\theta^2}{16\sqrt{2\pi}\sigma} \mathrm{e}^{-\frac{(\theta-\theta_0)^2}{2\sigma^2}} \mathrm{d}\theta, \end{aligned} \tag{3}$$

where d_L is the luminosity distance. Thus the number of bursts detected is,

$$N = \int \frac{1}{4}\pi n d_L^2 \mathrm{d}d_L \int_0^{(\frac{4E_0(1+z)}{\pi F d_L^2})^{\frac{1}{2}}} \frac{\theta^2}{\sqrt{2\pi}\sigma} \mathrm{e}^{-\frac{(\theta-\theta_0)^2}{2\sigma^2}} \mathrm{d}\theta. \tag{4}$$

Similarly, for an exponential distribution with the parameter λ,

$$N = \int \frac{1}{4}\pi n d_L^2 \mathrm{d}d_L \int_0^{(\frac{4E_0(1+z)}{\pi F d_L^2})^{\frac{1}{2}}} \theta^2 \lambda \mathrm{e}^{-\lambda\theta} \mathrm{d}\theta, \tag{5}$$

and for a power-law distribution with the parameter t,

$$N = \int \frac{1}{4}\pi n d_L^2 \mathrm{d}d_L \int_0^{(\frac{4E_0(1+z)}{\pi F d_L^2})^{\frac{1}{2}}} \theta^{2-t} \mathrm{d}\theta. \tag{6}$$

If we assume the peak luminosity of GRBs is the standard candle, we only need to replace E_0 and F by the standard candle luminosity L_0 and the peak flux P in equations (3), (4), (5) and (6), respectively. We number these slightly modified equations as (3)$'$, (4)$'$, (5)$'$ and (6)$'$, respectively; for brevity we do not list these equations explicitly.

3. Results

We apply equation (1) for the GRB space density distribution and equations (4), (4)$'$, (5), (5)$'$, (6) and (6)$'$ for the GRB opening angle distribution to fit the observed GRB intensity distribution. We focus our study on long GRBs ($T_{90} > 2s$) (Kouveliotou *et al.* 1993) because currently only long bursts are known to be originated at cosmological distances from the direct redshift measurements of the optical afterglows of some long GRBs. All long GRBs with peak flux values and detection probability above 5% included in the GRB 4B+ Catalog (Paciesas *et al.* 2000) are selected, resulting in 1727 GRBs as our sample. The fluence values of these GRBs are taken also directly from the catalog. For the peak flux distribution, the detection efficiency correction is done directly using the detection probability of each peak flux value given in the catalog. Because the fluence and peak flux does not have a good linear correlation, for each fluence range we average the detection probability of all GRBs included in this fluence range and then correct for the detection efficiency of this fluence range.

		Gaussian	**exponential**	**power-law**
Total energy as standard candle.	χ_r^2	281.3/194	**261.1/195**	623.2/195
		θ_0=0.028±0.007 σ=0.050±0.008	λ=**38.1±3.8**	t=4.51±0.09
	z_0	0.23 ± 0.07	**0.23±0.08**	0.27 ± 0.11
	z_{break}	$7.7 - \infty$	**7.9-∞**	$7.2 - \infty$
	n_0	231 ± 51	215 ± 45	207 ± 44
	E_0	0.69 ± 0.37	0.80 ± 0.41	0.72 ± 0.36

		Gaussian	**exponential**	**power-law**
Peak luminosity as standard candle.	χ_r^2	241.1/194	**227.1/195**	539.7/195
		θ_0=0.031±0.008 σ=0.047±0.01	λ=**34.4±2.1**	t=4.52±0.09
	z_0	0.23 ± 0.09	**0.24±0.07**	0.29 ± 0.09
	z_{break}	$9.5 - \infty$	**9.8-∞**	$9.2 - \infty$
	n_0	273 ± 44	264 ± 49	257 ± 39
	L_0	0.67 ± 0.32	0.64 ± 0.22	0.69 ± 0.30

Table 1: $\chi_r^2 = \chi^2/dof$. L_0 is in units of 10^{51} *ergs/s.* n_0 in units of Gpc^3/year. All errors quoted are for 68.3% confidence. For both cases the exponential distribution of the opening angle offers the best fits; however the residuals for the combination of peak luminosity as the standard candle and the exponential distribution is the only statistically acceptable one.



Fig. 1.— Comparison of different models with data. Top panel: long GRBs with fluence as standard candle; Lower panel: long GRBs with peak luminosity as standard candle.

Our fitting results are shown in *Table.1* and *fig.1.* Because our fittings are done with the integral distributions, the numbers of GRBs in different bins are not completely independent. To address this problem, we carried out extensive bootstrap simulations. Taking the 1727 GRBs as the seed distribution, we randomly generated a large numbers of peak flux or fluence distributions and fitted the sampled distributions in the same way. The best values and errors of all parameters are then calculated from these fittings; the bootstrapped parameter values and errors are consistent with that listed in Table 1. We therefore believe the correction between the different points does not have any significant impact to our results. From the fitting results, we conclude:

- For long GRBs, despite the different fitting residuals of the two standard candle models and three opening angle distributions assumed, the cosmological parameters of GRBs inferred are strikingly similar, indicating the robustness of the GRB cosmological model.
- The power-law distribution for the GRB opening angles cannot fit the data and is thus rejected confidently.
- The Gaussian distributions for the GRB opening angles fit the data only marginally, and may be rejected too.
- For both standard candle models the exponential distribution for the GRB opening angles provides the best fit to the data.
- The combination of the peak luminosity as standard candle and the exponential distribution for the GRB opening angles is the only one with statistically acceptable fit to the data; we therefore accept this as the accepted model for long GRBs. Under this model the peak luminosity of a GRB is $(0.64 \pm 0.22) \times 10^{51}$ *ergs/s*, the exponential constant of the opening angle distribution is 34 ± 2 (radian^{-1}), the lower bound to the GRB redshift is $z_0 = 0.24 \pm 0.07$, and the upper break redshift is $z_{break} > 9.8$ (no lower limit to the break is detected because not enough sufficiently faint GRBs have been observed with BATSE).

4. Discussion

First we compare our study with previous studies of GRB cosmological models with BATSE intensity distribution (Fenimore *et al.* 1993; Fenimore and Bloom 1995) (Wickramasinghe *et al.* 1993; Wijers *et al.* 1998). All these studies concluded that the dimmest GRBs are at redshift around unity, probaly following the measured star formation rate which peaks also at redshift around unity (Steidel *et al.* 1999). The main differences between our approach and previous studies are the following:

- We take the GRB opening angle distribution as part of the cosmological model. This is very important, since the observed GRB flux depends strongly on the opening angle, as well as the redshift.
- We separate the long and short GRBs as two different classes in our study. Our conclusions are only applicable to long GRBs. For short GRBs, no conclusion can be drawn due to the limited statistical quality of data.
- One simplification of our study is that we did not make the energy spectral correction to the fluence or peak flux as a function of z (K-correction); we just take the fluence or peak flux values as listed in the catalog. Technically this correction is not straightforward, because we can no longer uniquely convert the fluence or peak flux directly to redshift, because we believe that GRB radiation is beamed, and the opening angle follows a certain distribution. Therefore the observed fluence or peak flux is due to both the redshift and the opening angle of the GRB. The good fit of the assumed simple cosmological model with only several free-parameters indicates that this correction may not impact our conclusions significantly. In fact, since the energy spectra of GRBs in the BATSE band follows approximately a power-law with an index of -2, the K-correction is not important.

Frail *et al.* (2001) suggested that the total energy of GRBs may be a constant, based on the opening angles of several GRBs determined from their afterglow light curve break. Our result is not in major conflict with their suggestion, but favors the peak luminosity as the standard candle. Perhaps the direction of the GRB beam is not completely stable, as indicated from observations of other relativistic jets and outflows from stellar mass and supermassive black holes; only when the beam is pointed directly at us can we observe the maximum flux from the GRB. However it has been argued that for some GRBs the lower limit of the total energy released is 10^{52} *ergs* as determined from the redshifted X-ray emission lines (Ghisellini *et al.* 2002). However the reality of the detection of the emission lines is still under debate, and an alternative interpretation of the detected emission lines also exists (Wang *et al.* 2002).



Fig. 2.— The data for the obsered GRB opening angles are taken from Bloom *et al.* 2003; arrows indicate lower and upper limits of GRB opening angles. The power-law distribution fits the data for large opening angles, but fails to predict the absence of GRBs detected with opening angles smaller than 0.04 radian. The Gaussian distribution agrees with the peak reasonably well, but cannot predict enough GRBs with opening angles greater than 0.2 radian. The exponential distribution is consistent with the data.

The opening angles of some GRBs with afterglow detections have been determined previously from the GRB afterglow data (Frail *et al.* 2001; Bloom *et al.* 2003); we call this Frail-distribution. In *fig.2* we compare the Frail-distribution with the opening angle distributions determined with the observed GRB intensity distribution data, just as a sanity check. The power-law distribution agrees with the Frail-distribution for large opening angles, but fails to predict the absence of GRBs detected with opening angles smaller than 0.04 radian. No GRBs with opening angles narrower than 0.04 radian was directly observed, but the recent report on the polarization of GRB021206 suggests that it is possible for some GRBs to have opening angles narrower than 0.01 radian (Coburn and Boggs 2003; Waxman 2003). The Gaussian distribution agrees with the peak of the Frail-distribution distribution reasonably well; however this model cannot predict enough GRBs with opening angles greater than 0.2 radian. The exponential distribution is consistent with the Frail-distribution distribution for both standard candle models, but favors slightly the peak luminosity standard candle model. It was suggested that for the model in which the GRBs follow the star formation rate strictly, the most possible opening angle should be around 0.1 radian (Bloom *et al.* 2003). However, if the majority of faint GRBs originate at higher redshift than the peak of star formation rate as our model predicts, then a narrower average opening angle is required.

We therefore conclude that GRB opening angles follow an exponential distribution with a mean opening angle of $1/\lambda = 0.03$ radian, and the GRB space density follows a power-law with an index of -3.5 between $z_0 = 0.24 \pm 0.07$ and $z_{break} > 10$. The general trend of the GRB space density distribution determined here as a function of z is consistent with previous studies (Schaefer *et al.* 2001; Norris 2002); however our result is statistically much more robust, and in particular z_0 is determined for the first time and the lower limit to z_{break} is constrained more strictly than before. Our result therefore demonstrates that GRBs are no longer produced frequently in the nearby and present universe at $z < 0.24 \pm 0.07$, and GRBs are predominantly produced in the early universe at $z > 10$. Indeed most GRBs with direct redshift measurements have redshifts $z > 0.3$ (Djorgovski *et al.* 2001), with the exceptions of GRB 980425 (z=0.0085) and GRB 030329 (z=0.1687) which may be associated with two peculiar but similar supernovae SN1998bw (van Paradijs 1999), and SN2003dh (Stanek *et al.* 2003). It remains to be seen if these GRBs belong to a subset of GRBs which are different from the majority of GRBs which seem to be produced at much larger redshift.

It has been suggested that GRBs follow star formation rate, because it is commonly believed that GRBs are produced as the final gravitational collapse of massive stars (Djorgovski *et al.* 2001; Paczynski 1998). The GRB space density determined here follows the measured star formation rate between $z \approx 0.24$ to $z \approx 1$. The determination of $z_0 = 0.24 \pm 0.07$ suggests that the massive stars capable of producing the majority of long GRBs are no longer formed frequently today. Future GRB experiments capable of measuring the redshift of many more

GRBs and thus providing direct distance measurements of GRBs, such as the *Swift* mission to be launched in late 2003 (Gehrels 2000), will test this prediction firmly.

The monotonic increase of the GRB space density up to at least $z = 10$ suggests that the formation rate of massive stars capable of producing GRBs should peak at significantly higher z than the measured star formation rate. This implies that the "dark ages" of the universe ended much earlier than previously believed at $z \sim 4-5$ based on the measured space density distribution of high-redshift QSOs (Hook *et al.* 2002). There are also some previous independent works showing that an essential amount of long GRBs should be at $z > 4$ (Choudhury and Srianand 2002; Bagoly *et al.* 2003) or even up to $z \cong 20$(Meszaros and Meszaros 1996). A recently detected GRB031203 by INTEGRAL was reported to be located at $z > 9$ (formally $z \geq 11$) (GRBlog 2003), consistent with our model prediction. Therefore GRBs may be a probe of the epoch of reionization of the early universe (Lamb and Haiman 2003). Currently the GRB intensity distribution data cannot constrain the exact upper limit of z_{break}, because not enough extremely faint GRBs have been observed. Future GRB experiments, such as the proposed "Next Generation GRB Mission" (Grindlay *et al.* 2001) would detect many more GRBs originated in the extremely early universe, which may determine the exact ending time of the "dark ages" of the universe in order to test the prediction of some numerical simulations (Abel *et al.* 2002) which suggests that the first stars may have been formed at $z \sim 20$. GRBs may prove to be the most sensitive and ultimate probes into the early epoch of the universe when the first structures were formed.

Acknowledgement: We thank Drs. Zigao Dai, Tan Lu, Yongfeng Huang and Daming Wei for valuable comments to the manuscript. This study is supported in part by the Special Funds for Major State Basic Research Projects (10233010) and by the National Natural Science Foundation of China. SNZ also acknowledges supports by NASA's Marshall Space Flight Center and through NASA's Long Term Space Astrophysics Program.

REFERENCES

Abel, T., Bryan, G. L., & Norman, M. L. 2002 Science, 295, 93

Bagoly, Z., Csabai, A., Meszaros, A., Meszaros, P., Horvath, I., Balazs, L. G., & Vavrek, R. 2003, A&A, 398, 919

Balazs, L. G., Bagoly, Z., Horvath, I., Meszaros, A. & Meszaros, P. 2003, A&A, 401, 129

Bloom, J. S., Frail,D. A. & Kulkarni, S. R. 2003, ApJ, 594, 674

Choudhury, T. R, & Srianand, R. 2002, MNRAS, 336, L27

Coburn, w., & Boggs, S. E. 2003, Nature, 423, 415

Djorgovski, S., *et al.* 2001, *ESO Astrophysics Symposia, Berlin: Springer Verlag*

Fenimore, E.E., & Bloom, J.S. 1995, ApJ, 453, 25

Fenimore, E.E., *et al.* 1993, Nature, 366, 40

Frail, D. A. 2001, *et al.*, ApJ, 562, L55

Fruchter, A. S., *et al.*. 1999, ApJ, 519, L13

Gehrels, N. A. 2000, *Procspie* 4140, 42

Ghisellini, G., *et al.* 2002, A&A, 389, L33

http://grad40.as.utexas.edu/grblog.php?GCN=2468

Grindlay, J., *et al.* 2001, Presented at Woods Hole GRB Conf., to appear in AIP Conf. Proc.

Hook, I. M., *et al.* 2002, A&A, 391, 509

Kouveliotou, C., *et al.* 1993, ApJ, 413, L101

Lamb, D.Q., & Haiman, Z. 2003, AAS, HEAD2003, 16.15

Meegan, C. A., *et al.* 1992, Nature 355, 143

Meszaros, A., & Meszaros, P. 1996, ApJ, 466, 29

Norris, J. P. 2002, ApJ, 579, 386

Paciesas, W. S., *et al.*, *VizieR On-line Data Catalog: IX/20A.* Originally published in: 2000, ApJ, 122, 465P; ApJ, 122, 497H

Paczynsky, B. 1998, ApJ, 494, L45

Schaefer, B. E., Deng, M. & Band, D.L. 2001, ApJ, 563, L123

Stanek, K.Z., *et al.* 2003, ApJ, 591, L17

Steidel, C. C., *et al.* 1999, ApJ, 519, 1

van Paradijs, J. 1999, Science, 286, 693

Wang, W., Zhao, Y., & You, J. H. 2002, ApJ, 576, L37

Waxman, E., Kulkarni, S. R. & Frail, D. A. 1998, ApJ, 497, 288

Waxman, E. 2003, Nature, 423, 388

Wijers, R.A.M.J., *et al.* 1998, MNRAS, 294, L13

Wickramasinghe, W.A.D.T., *et al.* 1993, ApJ, 411, L55

Woosley, S. 2001, Nature, 414, 853

Correspondence should be addressed to S.N.Z. (e-mail: zhangsn@tsinghua.edu.cn).

This preprint was prepared with the AAS LaTeX macros v5.2.